

## RESEARCH ARTICLE

### The effect of Azolla extract on vegetative growth indicators of three varieties of Barley (*Hordeum vulgare* L.)

Rahim B. Al-Nasr-Allah <sup>1\*</sup> and Ali R. Alhasany <sup>1</sup>

<sup>1</sup> Department of Field Crop, College of Agriculture, University of Al-Muthanna, 62001, Iraq

#### ABSTRACT

The field trial was executed on land belonging to a planter in the Al Bandar area (3 km from the center of Samawah) during the 2022-2023 season to determine the effect of four levels of Azolla extract (45, 30, 15, 0 liters ha<sup>-1</sup>) on three varieties. From barley (AXAD, Ibaa 99, Ibaa 265), the trial was conducted utilizing a block-design randomized completely according to the split-plot method within tri-replicates, with the varieties occupying the main panels and the Azolla levels in the secondary panels. The results exhibited that the grade of 30 liters of ha<sup>-1</sup> was significantly outstanding to the flag leaf area, the flag leaf content of chlorophyll, the length of the spike, and the number of branches per square meter, as it gave averages of (24.48 cm<sup>2</sup>), (44.63 SPAD), (5.82 cm), and (509.40 beches m). 2) Sequentially. The results also exhibited that the Aksad variety was superior in the nitrogen content of the plant, the flag leaf's chlorophyll content, the spike-length and the branches-number per square meter, and gave averages of (2.47%), (45.22 SPAD), (6.17 cm), and (512.50 Shatam-2), respectively, and the variety was superior. Fathers 265 in plant height and leaf area, and gave two averages of (104.05 cm) and (23.33 cm<sup>2</sup>), respectively.

**Keywords:** Barley; Azolla; vegetative growth indicators

#### Corresponding Author:

Rahim. B. Al-Nasr-Allah, Department of Field Crop, College of Agriculture, University of Al-Muthanna, 62001, Iraq.

E-mail: [rheem5846@gmail.com](mailto:rheem5846@gmail.com)

Received: 10/12/2024

Accepted: 18/02/2025

Published: 31/03/2025

DOI: <https://doi.org/10.64333/MJAE.25.1.4>

Copyright © 2025. This is an open-access article distributed under the Creative Commons Attribution License.

## INTRODUCTION

The barley crop is considered an important strategic grain crop. It is one of the productive plants that were domesticated in ancient times, and it contributes mainly to achieving human food security. It ranks fourth in importance after wheat, corn, and rice. It is the fundamental exporter of bread in the obsolete scientist, as its grains are opulent in proteins and carbohydrates essential for human nutrition, as well as a altitude level of fiber, oil, and vitamin B (Gani and Salman, 2011). Due to the lack of production in the barley crop, there was a need to increase the production of varieties with good genetic characteristics and high production that are suitable for the environmental conditions in the region, as barley varieties vary in their production capacity depending on the variation in their genetic makeup and the extent of their influence on environmental conditions on the one hand, and the lack of control over production techniques on the other. For this reason, many researchers were interested in searching for new ways to raise productivity while improving quality by exploiting the biodiversity found in varieties and genetic compositions with high yields (Al-Sahuki, 2006; Al-Nasr-Allah and Alhasany, 2023). Azolla extract is one of the important organic nutrients because it contains nutrients, amino acids, and growth regulators that have an important role in physiological processes as well as improving the growth and production of plants

(Altai *et al.*, 2019). The Azolla plant is a fern that reproduces sexually and asexually and belongs to the Azollaceae family. Which lives on the surfaces of drains and stagnant water and is characterized by rapid growth and spread, as its biomass can be doubled significantly every 2-3 days (Cohen *et al.*, 2002).

Azolla contains 3-4% dry matter and is considered a rich source of fundamental amino-acids, and A, and B12 vitamins. It comprises many elements such as calcium, phosphorus, potassium, iron, magnesium, and copper. Blue-green algae (*Anabaena azolla*) that coexist symbiotically with Azolla fixes nitrogen. At a higher rate than legumes, ranging between 100-170 kg nitrogen ha<sup>-1</sup>, Azolla is a rich source of protein, as it constitutes 25-30% protein, minerals 15%, and more than 7% a mixture of amino acids, biologically active materials, and polymers (Pillai *et al.*, 2005). Our study aims to determine the best combination of interaction between varieties and Azolla extract, which gives the best prosperity and productivity barley crops.

### MATERIALS AND METHODS

The trial was executed during the period (2022-2023) in Al-Bandar district fields, southwest of Al-Muthanna Governorate (3 km from Samawah), to study the role of Azolla extract at four levels (0, 15, 30, and 45 liters) on the prosperity and productivity of three barley varieties (Aksad, Abaa 99, Abaa 265). Ploughing, leveling operations, and dividing were executed, and the trial field was planned within the design utilized. Tri-varieties seeds were sown on November 3 (Al-Jayashi *et al.*, 2020). The trial soil was randomly distributed into 36 trial units. The area of the experimental unit was 4 m2 (2 x 2 m), containing 10 lines between two lines and the last 20 cm, and a seed rate of 120 kg ha-1 was adopted (General Authority for Agricultural Research, 2011). The fertilization-nitrogen operation was executed by adding about 120 kg ha-1 [(CO(NH<sub>2</sub>)<sub>2</sub> 46% N)] in 2 batches, and it was sufficient to add the first batch only after 15 days of planting, after calculating the amount of nitrogen present in the soil (Table 1). The fertilization-phosphate operation was carried out by adding about 100 kg ha-1 [Ca(H<sub>2</sub>PO<sub>4</sub>)<sub>2</sub>·H<sub>2</sub>O) P<sub>2</sub>O<sub>5</sub> 46%] in 1 batch at sowing as described by Al-Abedi (2011). Fertilizer-potassium was applied about 80 kg ha-1 in the form of potassium sulphate [K<sub>2</sub>SO<sub>4</sub> (K<sub>2</sub>O 50%)] in 1 batch at sowing (Ali *et al.*, 2014). Irrigation and weeding were also carried out whenever necessary. The Azolla plant was grown in the home garden's 4 m cement basin. A mixture of clay and cow manure was placed inside the basin, and then fresh water was added from the tap, and then placed 2 kilograms of live Azolla plants obtained from one of the farms that produced it. The Azolla plant needed 3-4 days to double its biomass. The Azolla was harvested to make the extract after 7 days. Random samples were taken from various venues of every replicate and mixed to take a composite specimen representing the experimental field and from depth (0-30), and some physical and chemical analyzes of the experimental field were conducted on it before planting, which are shown in Table 1.

**Table (1)** The physicochemical characteristics of the experimental field pre-planting

Units	Values	Attribute	
	7.20	pH	
ds/m <sup>-1</sup>	4.50	E.c (1:1)	
Mg/kg dry weight	26.60	Nitrogen(available)	
Mg/kg dry weight	13.20	phosphorus(available)	
Mg/kg dry weight clay	118.70	potassium(available)	
	22.33	Clay	
g/kg <sup>-1</sup>	29.12	Silt	Soil separators
	48.55	Sand	
Sandy loam		Soil texture	

## **Studied attributes**

### **1- Nitrogen concentration in plants (%)**

Samples were collected from the shoots of each trial unit at the 75% flowering stage, and the nitrogen concentration was calculated according to the method described by [Cresser and Parsons \(1979\)](#) in a nitrogen distillation device (Macrokjeldhal).

### **2- Plant height (cm)**

It was measured using a ruler from the soil surface to the end of the spike, without cutting, for 10-plants collected randomly for each trial unit at harvest.

### **3- Area of the flag leaf (cm<sup>2</sup>)**

It was calculated at the stage of completion of flowering, and 10-plants were randomly collected from the midlines of each trial unit, according to the formula listed in [Thomas \(1975\)](#).

(Area of flag paper = length of paper x maximum width x 0.75).

### **4- Chlorophyll (SPAD) content of leaves**

The total chlorophyll content of leaves was estimated using the SPAD Chlorophyll Meters (Spectrum Technologies Inc.) as an average of ten plants, taken randomly from the midlines of each experimental unit at the stage of complete flowering.

### **5- Spike length (cm)**

It was calculated as an average of ten spikelets taken randomly from the harvested midlines. The length of the spike was calculated from the basal of spikelet-terminal to the top of the upper terminal spikelet.

### **6-Tillers-number per square meter (tillers m<sup>-2</sup>)**

The numbers of tillers were calculated from harvesting two middle lines in the fully mature stage, and the harvested area represented 80 cm, which was then converted to square metres.

## **RESULTS AND DISCUSSION**

### **1- Nitrogen concentration in plants (%)**

The results of Table (2) indicated the superiority of the fourth level, A3, which gave an average of (2.72%) and didn't vary incomparably from the third level, A2, which was a moderate (2.71%). They vary extremely from the comparison treatment A0, which gave a moderate of (1.64%). The reason for increasing the nitrogen content by increasing the levels of addition may be due to the Azolla extract's content of nutrients and amino acids, which helped the plant to grow in general, and thus this is reflected in good root growth when some amino acids are available. This helps directly in stimulating the roots, such as the amino acid methionine, which contributed to absorbing the largest quantities of element-nitrogen, thus increasing the percentage of parts-plant. This is consistent with [\(Al-Jabri and Al- hasany, 2022\)](#), as he indicated an increase in the concentration of nitrogen in the plant. As shown, the data Table (2) indicates the superiority of the Aksad variety, as it gave an average of (2.47%) and didn't vary incomparably from variety Aba 265, which gave a moderate of (2.27%). They differed significantly from the variety Aba 99, which recorded the lowest average of (1.90%). The reason may be due to the difference. The genetic composition of the varieties in the nature of root growth and their ability to absorb nitrogen from the soil, as well as the nature of the vegetative growth of the varieties and their nutritional requirements, and this is consistent with [\(Al-Hassany, 2021\)](#), as he indicated the difference in the concentration of nitrogen in the plant according to the varieties.

The data in Table (2) indicated that the interaction was significant between the Azolla extract and the varieties, as the fourth level of extract A3 with the variety Oxad, the combination ( $V1 \times A3$ ), shown the highest, a moderate, amounting to (2.92%), while the combination represented by the second level of extract A1 with the variety Aba gave 99 ( $V2 \times A1$ ) The lowest average was (1.29%).

**Table (2)** The impact of Azolla extract, varieties, and interaction among on plant nitrogen concentration (%)

Azolla Cultivars	Axad	IPA99	IPA265	Azolla average
A0	1.94	1.51	1.48	1.64
A1	2.16	1.29	1.94	1.80
A2	2.86	2.48	2.78	2.71
A3	2.92	2.34	2.89	2.72
Cultivars average	2.47	1.90	2.27	
<b>L.S.D</b> <b>(0.05)</b>	<b>Cultivars</b> 0.24	<b>Azolla</b> 0.18	<b>Interaction</b> 0.32	

## 2- Plant height (cm)

The results of Table (3) indicated that the comparison treatment A0 was significantly superior and gave an average of (104.48 cm), with vary extremely from the third and fourth levels, A2 and A3, which gave averages of (100.33 and 100.32 cm), respectively. This may be due to the role of some amino acids in Azolla extract helps in early crop growth, thus prompting the plant to flower and not wasting nutrients manufactured in increasing plant height, such as the amino acids isoleucine, lysine, and glutamic acid. It was renowned in the data of Table (3) that there was an extreme superiority for the variety Aba 265, as it gave an average of 104.05 cm. It didn't vary incomparably from the variety Aba 99, which gave a moderate of 102.35 cm. It differed significantly from the variety Aksad, which exhibited the lowest average of 98.62 cm. The variation of varieties in plant height is due to the difference in genetic composition between the varieties, which is in agreement with [Al-Hassany \(2021\)](#). The data of Table (3) also showed that the interaction is greatly between the Azolla extract and the varieties, as the first level A0 with the variety Aba 265, the combination ( $V3 \times A0$ ), shown the highest, a moderate, amounting to (107.72 cm), whereas the integration represented by the fourth level A3 and the variety Kassad ( $V1 \times A3$ ) had the lowest average (97.85 cm).

**Table (3)** The impact of Azolla extract, varieties, and interaction among on plant height (cm)

Azolla Cultivars	Axad	IPA99	IPA265	Azolla average
A0	99.78	105.93	107.72	104.48
A1	98.85	102.00	103.83	101.56
A2	98.00	100.80	102.20	100.33
A3	97.85	100.68	102.43	100.32
Cultivars average	98.62	102.35	104.05	
<b>L.S.D</b> <b>(0.05)</b>	<b>Cultivars</b> 3.82	<b>Azolla</b> 0.89	<b>Interaction</b> 3.74	

## 3- Area of the flag leaf (cm<sup>2</sup>)

The data of Table (4) indicated the superiority of the third level, A2, as it gave an average of (24.48 cm<sup>2</sup>) and didn't vary extremely from the fourth level, A3, which show a moderate of (23.30 cm<sup>2</sup>), and they differed significantly with the comparison treatment A0, which displays the lowest an moderate of (19.06 cm<sup>2</sup>). The area of flag leaves is increased when treated with Azolla extract may be due to the nutritional elements and amino acids it contains, which help in increasing activities of vital functions, and the increment in the nitrogen content of the leaves (Table 2), which increases the process of

division and elongation-cell and thus increment the flag leaf area, consistent with (Saudi, 2017).

The data in Table (4) indicated the superiority-variety Aba 265, as it show a moderate of (23.33 cm<sup>2</sup>) and didn't vary extremely from the variety Axad, which show a moderate of (22.03 cm<sup>2</sup>), and they varied extremely from the variety Aba 99, which show a moderate (20.13 cm<sup>2</sup>). The superiority of the cultivar Abaa 265 in flag leaf area is due to the difference in its genetic composition from the rest of the varieties under study. The increment the flag leaf area in the cultivars Abaa 265 and Aksad may be due to an increase in the nitrogen content of the plant (Table 2), which helped in increasing the vital processes in the plant and was reflected positively in increasing the area of the flag leaf, which is consistent with Al-Jayashi *et al.*, (2020). The data in Table (4) also indicated that the synergy was extremely strong between the Azolla extract and the cultivars, as the third level of the extract A2 with the cultivar Aba 265, the combination (V3 × A2), gave the highest average for this trait, amounting to (25.71 cm<sup>2</sup>), and it did not differ significantly from many of the combinations, while the comparison treatment gave A0 with variety Aba 99 (V2 × A0) had a lower average of (17.98 cm<sup>2</sup>). This may be explained by what was discussed in the effect of individual factors and this was reflected in the interaction.

**Table (4)** The impact of Azolla extract, varieties, and interaction among on flag leaf area (cm<sup>2</sup>).

<b>Azolla Cultivars</b>	<b>Axad</b>	<b>IPA99</b>	<b>IPA265</b>	<b>Azolla average</b>
A0	18.56	17.98	20.63	19.06
A1	18.66	20.70	22.09	20.48
A2	25.39	22.34	25.71	24.48
A3	25.52	19.48	24.90	23.30
Cultivars average	22.03	20.13	23.33	
<b>L.S.D (0.05)</b>	<b>Cultivars 1.58</b>	<b>Azolla 1.66</b>	<b>Interaction</b>	<b>2.73</b>

#### 4- Science leaf content of chlorophyll (SPAD)

The data of Table (5) that there was an extremely superiority for level A2, as it gave an average of (44.63 SPAD) for this characteristic, and it didn't vary extremely with levels A3 and A1, which gave averages of (44.08 and 43.02 SPAD), respectively, and it differed significantly with the treatment A0. It gave the lowest average (41.72 SPAD). The flag leaf 'are increasing by chlorophyll content to the nutritional elements and amino acids contained in the Azolla extract, as there is an important role for some amino acids that contribute to the formation of the chlorophyll molecule, such as the amino acid alanine, which contributed In increasing the leaf content of chlorophyll, which is consistent with AL-Hasany *et al.*, (2019). The data in Table (5) indicated that the Aksad variety was significantly superior, as it gave an average of (45.22 SPAD) and didn't vary extremely with the variety Abaa 265, which recorded an average of (43.45 SPAD), but it differed significantly with the variety Abaa 99, which displays the lowest an moderate of (41.43 SPAD). The chlorophyll content discrepancy in the flag leaf is explained by the genetic nature of the Aksad variety and the amplitude of its response to the surrounding ecological conditions better than the rest of the varieties included in the study, and this is consistent with, and this is consistent with (Al-Jayashi *et al.*, 2020).

**Table (5).** The impact of Azolla extract, varieties, and interaction among on the chlorophyll content of the flag leaf (SPAD)

Azolla Cultivars	Axad	IPA99	IPA265	Azolla average
A0	43.25	40.03	41.89	41.72
A1	44.93	40.64	43.50	43.02
A2	46.35	42.57	44.98	44.63
A3	46.34	42.47	43.43	44.08
Cultivars average	45.22	41.43	43.45	
<b>L.S.D (0.05)</b>	<b>Cultivars</b>	<b>Azolla</b>	<b>Interaction</b>	
	2.65	1.68	N.S	

### 5- Spike length (cm)

The data of Table (6) exhibited an extremely superiority for the third level, A2, as it show a moderate of (5.82 cm). It didn't vary extremely with the fourth level, A3, which show a moderate of (5.42 cm). They differed significantly with the comparison treatment A0, which displays the lowest a moderate of (5.11 cm). The length of the spike is increasing due to the increment in the flag leaf area, Table (4), resulting from increasing the levels of addition to Azolla extract, thus increasing carbon metabolism and providing the largest amount of nutrients, which led to an increment in the length of the spike, which is consistent with [Altai et al., \(2019\)](#), as They indicated that the length-spike was increased by adding plant extracts.

The data in Table (6) showed that there was a notably superiority of the Aksad variety in spike length, as it gave an average of (6.17 cm), and a notably vary with the two varieties Aba 99 and Aba 265, which gave averages of (5.05 and 5.03 cm), respectively. This discrepancy between the varieties may be due to length, where spike is due to the difference in genetic composition between them, and this is consistent with [Al-Hassany \(2021\)](#), who showed the variation in spike length according to the variety.

**Table (6)** The impact of Azolla extract, varieties and and interaction among on spike length (cm).

Azolla Cultivars	Axad	IPA99	IPA265	Azolla average
A0	5.66	4.62	5.07	5.11
A1	6.41	4.92	4.64	5.32
A2	6.44	5.17	5.84	5.82
A3	6.18	5.49	4.58	5.42
Cultivars average	6.17	5.05	5.03	
<b>L.S.D (0.05)</b>	<b>Cultivars</b>	<b>Azolla</b>	<b>Interaction</b>	
	0.50	0.44	N.S	

### 6- Number of beaches per square meter (tillers M<sup>-2</sup>)

The data in Table (7) showed a significant superiority of the third level of Azolla A2, which gave an average of (509.40 tillers m<sup>-2</sup>) for this characteristic, and didn't vary extremely from the fourth level, A3, which show a moderate of (495.60 tillers m<sup>-2</sup>), and they differed significantly with the treatment A0, which displays the lowest an moderate was (451.40 tillers m<sup>-2</sup>). The number of strands is increasing when adding the Azolla extract due to the extract's content of the growth regulator kinetin, which is a cytokinin that causes breaking the apical dominance and encouraging lateral stranding, which led to an increase in the number of tillers per square meter. This is consistent with [Al-Ghazal et al. \(2021\)](#), who indicated that the number of fractions increases with increasing concentrations of the additive. The data of Table (7) indicated that the Aksad variety was extremely superior and gave an average number of tillers that amounted to (512.50 tillers m<sup>-2</sup>) and didn't vary extremely from the variety Ibaa 265, which show a moderate of (501.60 tillers m<sup>-2</sup>), and they vary extremely with the variety Ibaa 99, which displays the lowest an moderate.

Owing to this characteristic, it accomplished (438.30 tillers m<sup>-2</sup>), The differences among the varieties of number-tillers may be due to their differences in their genetic composition and the extent of the variety's ability to beach under the environmental conditions of the region, and this is consistent with (Al-Absawi *et al.*, 2023).

**Table (7)** The effect of Azolla extract, varieties, and the interaction between them on the number of trees per square meter (tillers m<sup>-2</sup>)

<b>Azolla Cultivars</b>	<b>Axad</b>	<b>IPA99</b>	<b>IPA265</b>	<b>Azolla average</b>
A0	508.50	403.90	441.70	451.40
A1	488.20	450.40	501.70	480.10
A2	514.70	463.40	550.00	509.40
A3	538.60	435.40	512.90	495.60
Cultivars average	512.50	438.30	501.60	
<b>L.S.D (0.05)</b>	<b>Cultivars</b> 59.62	<b>Azolla</b> 33.94	<b>Interaction</b> N.S	

## CONCLUSIONS

The 30 L/ha level of Azolla extract significantly positively affected key barley traits, including flag leaf area, chlorophyll content, spike length, and the branches-number per m<sup>2</sup>, making it the most effective orientation. The Aksad variety outperformed the others in nitrogen content; chlorophyll content, spike length, and tillers density, indicating its strong response to Azolla extract application. The Ibaa 265 variety are excelled in plant height and leaf area, suggesting its suitability for conditions requiring taller plants with a larger photosynthetic surface. Azolla extract can be a natural bio-fertilizer to improve barley growth and productivity, reducing reliance on chemical fertilizers and promoting the environment.

**Disclaimer:** None

**Conflict of interest:** The authors declare that they have no relevant financial or non-financial interest to disclose.

**Funding disclosure:** None

## REFERENCES

- Gani, A. J. and K. A. Salman (2011). Barley from agriculture to harvesting. Published by The General Authority for Agricultural Research.
- Al-Sahuki, M. (2006). Plant breeding with the help of molecular parameters. Iraqi Agricultural Sciences Journal 37(4):67-72.
- Al-Nasr-Allah, R. B. and Alhasany A. R. (2023). Effect of Azolla Extract on Yield, Its Components and Protein Content in three Cultivars of Barley. Al-Muthanna Journal of Agricultural Sciences,2(10):1-13.
- Altai, D. S., Alhasany, A. R., Noaema, A. H., Idan, W. J., and Al-Farhan, I. M. (2019). Effectiveness of Azolla Extract In Reducing Use of Mineral Fertilizers (NPK) and Increasing Productivity of Wheat under New Reclaimed Soils. Journal of Plant Production, 10(10). 859-866.
- Cohen, M. F., Meziane, T., Tsuchiya, M., and Yamasaki, H. (2002). Feeding deterrence of Azolla in relation to deoxyanthocyanin and fatty acid composition. Aquatic Botany, 74(2), 181-187.
- Pillai, K. P., Premalatha, S. and Rajamony, S. (2005). Azolla-A sustainable feed substitute for livestock available at leisa magazine :26-27.
- Al-Jayashi, M. T. A. (2020). The effect of mowing, planting dates, and varieties on the growth characteristics and yield of fodder and grains of barley (*Hordeum vulgare* L.). Doctoral dissertation. Faculty of Agriculture. Al-Muthanna University.
- General Authority for Agricultural Research (2011). Barley from planting to harvest. Ministry Of Agriculture. General Authority for Agricultural Guidance and Cooperation. The Republic of Iraq.
- Al-Abedi, J. I. (2011). Guide to the use of chemical and organic fertilizers in Iraq. General Authority for Agricultural Guidance and

Cooperation. Ministry Of Agriculture. Iraq.

- Ali, N. S., Hamdallah S. R., and Abdel-Wahab A. S. (2014). Soil fertility. Ministry of Higher Education and Scientific Research - University of Baghdad. First edition. Dar Al-Kutub for Printing and Publishing.
- Cresser, M. S. and Parsons, J. W.( 1979). Sulphuric- perchloric acid digestion of plant material for the determination of nitrogen, phosphorus, potassium, calcium and magnesium. *Analytica Chimica Acta*, 109: 431-436.
- Thomas, H. (1975). The growth responses to weather of simulated vegetative swards of a single genotype of *Lolium perenne*. *The Journal of Agriculture Sciences*, 84(2), 333-343
- Al-Jabri , A. F. and Al- hasany. A. R. (2022). The role of fractionation of humic acids and spraying with Azolla extract on vegetative growth indicators growth of sunflower crop (*Helianthus annuus* L.). *Al-Muthanna Journal of Agricultural Sciences*, 9(2):1-8.
- Al-Hassany, R. T.J. (2021). The effect of nano- and conventional NPK fertilizer on the growth and yield of three varieties of barley (*Hordeum vulgare* L.). Doctoral dissertation. Faculty of Agriculture. Al-Muthanna University.
- Saudi, A. H. (2017). The effect of foliar spraying with seaweed extract on the growth, yield, and seed strength of bread wheat varieties (*Triticum aestivum* L.). *Iraqi Agricultural Sciences Journal*. (5)48: 1313-1325.
- AL-Hasany, A. R. K; Aljaberi M. A. R. and Alhilfi, S. K. J. (2019). Effect of Spraying with Seaweed Extract on Growth and Yield of Two Varieties of Wheat (*Triticum aestivum* L.). *Basrah Journal of Agricultural Sciences*, 32: 124-133.
- Al-Ghazal, S. A. Y., Maysar, M. A., and Al-Juhaishi, W. S. (2021). Response of growth and yield of sorghum (*Sorghum bicolor* L.) to different levels of humic acid and seaweed extract. *Al-Muthanna Journal of Agricultural Sciences*. (4)8 11-1.
- Al-Absawi, M. R. A. (2023). The effect of coating urea with white cement on the growth characteristics and yield of fodder barley (*Hordeum vulgare* L.) and soft wheat (*Triticum aestivum* L.). Doctoral dissertation. Faculty of Agriculture. Al-Muthanna University